arXiv:0908.1551v1 [math.PR] 11 Aug 2009

The two-type continuum Richardson model: Non-dependence of the survival of both types on the initial configuration

Sebastian Carstens [1] and Thomas Richthammer [2]

Abstract

We consider the model of Deijfen et al. for the competing growth of two infection types in $\mathbb{R}^d$, based on the Richardson model on $\mathbb{Z}^d$. Stochastic ball-shaped infection outbursts transmit the infection type of the center of the ball to all points of the ball that are not yet infected. Relevant parameters of the model are the initial infection configuration, the (type-dependent) growth rates and the radius distribution of the infection outbursts. The main question is that of coexistence: For what values of the parameters is there a positive probability that both types grow unboundedly? It is known that for this question the initial configuration basically is irrelevant, provided certain technical assumptions on the radius distribution are satisfied. Here we show how to get rid of these assumptions, introducing a slight generalization of the model, where immune regions and delayed initial infection configurations are allowed.

Key words: Continuum growth model, Richardson's model, initial configuration, competing growth, shape theorem.

Classification: Primary 60K35, Secondary 82B43.

1 Introduction

We consider the competing growth of n types of infections on an underlying space ($\mathbb{Z}^d$ in a discrete setting and $\mathbb{R}^d$ in a continuum setting, where $d \geq 1$ is the number of spatial dimensions). Initially, given (disjoint) bounded regions Γ_i are known to be i-infected, i.e. infected with type i. Whenever a region is i-infected, it will stay like that forever, and will try to i-infect healthy "neighboring" regions by means of stochastic infection outbursts that occur at a rate proportional to a given growth parameter β_i. A situation like that can be modelled by a stochastic process $Z = (Z_t^i)_{i\in\{1,\dots,n\},t\geq 0}$, where Z_t^i denotes the subset of the underlying space that is infected with type i at time t, such that for fixed i the process $(Z_t^i)_t$ is increasing, and for fixed t the sets $(Z_t^i)_i$ are disjoint. In the

---

[1] Mathematisches Institut der LMU, Theresienstr. 39, 80333 Muenchen, Germany

[2] Department of Mathematics, UCLA, Los Angeles, CA 90095-1555, United States

Research supported in part by NSF grants DMS-0300672 and DMS-0301795

one-type model, it is an interesting problem to investigate the asymptotic shape of Z_t/t for $t \to \infty$, i.e. the limiting shape of the rescaled infected area. (Here in the discrete setting every point of $\mathbb{Z}^d$ in Z_t should be replaced by the unit square centered at this point to make this meaningful.) In a multi-type model another interesting question is that of coexistence: Does it happen with positive probability that each of the infection types grows unboundedly, i.e. infects regions arbitrarily far away from the origin? The answer to both problems will at least to some extent depend on the parameters of the model considered, i.e. on the initial configuration $\Gamma = (\Gamma_1, \ldots, \Gamma_n)$ and the growth rates $(\beta_1, \ldots, \beta_n)$.

In 1973 D. Richardson introduced the first model of the above type in [R], considering a one-type model in $\mathbb{Z}^d$. This model was generalized to two types by O. Häggström and R. Pemantle in [HP1]. In these models the dynamics of infection outbursts can be described as follows: Every non-infected site gets infected at a rate of β_i times the number of i-infected nearest neighbors. The original paper [R] proved the existence of a non-random asymptotic shape in the one-type model. This was improved and generalized later on, and while it was shown that the asymptotic shape is convex and compact, not much is known about its quantitative features. In the two-type model O. Häggström and R. Pemantle showed that coexistence is possible, provided $\beta_1 = \beta_2$, see [HP1]. They conjectured that for $\beta_1 \neq \beta_2$ coexistence is not possible, and the best result in this direction so far is that for fixed β_1 there can be at most countably many values of β_2 such that coexistence is possible, see [HP2]. Both results assume an initial configuration consisting of two points, but in [DH2] M. Deijfen and O. Häggström showed that the question of coexistence basically is independent of the initial configuration (as long as in the initial configuration no type strangles the other in that it surrounds the initial configuration of the other type completely).

The situation for the corresponding model on $\mathbb{R}^d$ is very similar. The one-type continuum growth model was introduced by M. Deijfen in [D], and the first two-type version of this by M. Deijfen, O. Häggström and J. Bagley in [DHB]. For describing the dynamics of the infection outbursts in these continuum models, we fix a time t. The waiting time for the first outburst of infection type i after time t is exponentially distributed with rate $\beta_i \lambda^d(Z_t^i)$, where λ^d denotes Lebesgue measure. The outburst has the shape of a ball, where the center is chosen uniformly in Z_t^i and the radius is chosen w.r.t. a given radius distribution ρ. ρ is a further parameter of the model, and usually results rely on assumptions on the tail of ρ, namely

$$\int_0^\infty r^d d\rho(r) < \infty \quad \text{or} \tag{1.1}$$

$$\int_0^\infty e^{\varphi r} d\rho(r) < \infty \quad \text{for some} \quad \varphi > 0, \tag{1.2}$$

and conditions on the support of ρ such as

$$\rho((0,\delta)) > 0 \quad \text{for all} \quad \delta > 0. \tag{1.3}$$

We will assume (1.1) throughout this article, as this condition ensures that the model is well defined, and we note that (1.2) implies (1.1). In the one-type continuum model the asymptotic shape is known to be a ball, see the original article by M. Deijfen [D] and the generalization by M. Deijfen, O. Häggström and J. Bagley in [DHB], where the condition on ρ is (1.2); we have stated this result as Theorem 3 in Section 2.4. The results concerning coexistence are almost completely analogous to those in the discrete case: For $\beta_1 = \beta_2$ coexistence is possible, see M. Deijfen and O. Häggström in [DH1], and for fixed β_1 at most countably many values of β_2 allow coexistence, see [DHB]. Both results assume condition (1.2) and concern models with special initial configurations (consisting of two disjoint balls). But again the question of coexistence is basically independent of the initial configuration, see [DHB]. The last result assumes conditions (1.2) and (1.3) on ρ, and the aim of the present article is to show how the technical assumption (1.3) can be avoided. (In fact it will turn out that assuming condition (1.1) will be sufficient.) As an immediate consequence the aforementioned coexistence results extend to basically all initial configurations for all radius distributions satisfying (1.2). This is desirable as there are interesting radius distributions satisfying (1.2), but not (1.3), e.g. a degenerate ρ (i.e. $\rho = \delta_r$ for some $r > 0$), which gives a continuum growth model, where all outbursts have the same deterministic radius.

From now on we will refer to the growth model on $\mathbb{R}^d$ as described above as (d-dimensional n-type) continuum Richardson model (CR-model). For stating our result let us consider a two-type CR-model with initial condition $\Gamma = (\Gamma_1, \Gamma_2)$ and growth rates (β_1, β_2). Let B_Γ denote the smallest ball centered at the origin containing $\Gamma_\cup := \Gamma_1 \cup \Gamma_2$. Let L_i be the event that type i leaves B_Γ, G_i the event that type i reaches points arbitrarily far from the origin, and $G := G_1 \cap G_2$ the event of unbounded growth of both types. Considering more than one model we will indicate the relevant parameters in parentheses after the corresponding event.

Theorem 1 *We consider two d-dimensional ($d \geq 2$) two-type CR-models with initial configurations Γ and Γ' respectively, growth rates $\beta_i = \beta_i'$ and radius distributions $\rho = \rho'$ satisfying* (1.1)*. If we have $\mathbb{P}(L_i(\Gamma)) > 0$ and $\mathbb{P}(L_i(\Gamma')) > 0$ for $i = 1, 2$, the possibility of coexistence doesn't depend on the initial condition:*

$$\mathbb{P}(G(\Gamma)) > 0 \quad \Longleftrightarrow \quad \mathbb{P}(G(\Gamma')) > 0.$$

We note that the condition $\mathbb{P}(L_i(\Gamma)) > 0$ and $\mathbb{P}(L_i(\Gamma')) > 0$ for $i = 1, 2$ just states that in the initial configurations no type strangles the other, and condition (1.1) is assumed only because we need the models to be well defined. As mentioned above, combining Theorem 1 with the coexistence results from [DH1] and [DHB] immediately gives the following result:

Theorem 2 *We consider a d-dimensional ($d \geq 2$) two-type CR-model with initial configuration Γ, growth rates $\beta = (\beta_1, \beta_2)$ and radius distribution satisfying (1.2).*

(a) If $\beta_1 = \beta_2 > 0$ and $\mathbb{P}(L_i(\beta)) > 0$ for $i = 1, 2$ we have $\mathbb{P}(G(\beta)) > 0$.

(b) For fixed β_1 we have $\mathbb{P}(G(\beta)) = 0$ for all but countably many values of β_2.

The fact that in Theorem 1 condition (1.2) can be relaxed to condition (1.1) is because the proof does not rely on the full assertion of the Shape theorem, but only on the lower bound on the asymptotic shape. If ρ has more weight on larger values of the radius, the intuition is that the infected region grows even faster, so the lower bound on the asymptotic shape is still valid.

To give some idea on how to get rid of condition (1.3), we will give a short review of the main idea of the corresponding result in [DHB]: Let the two growth models with the given initial configurations be denoted by Z and Z'. We know that in Z coexistence is possible, and want to show the same for Z'. Assume that we can find balls $B_1 \subset B_2$ and a time T such that B_1 contains both initial configurations, in the model Z all regions infected at time T are within B_2 and all outbursts in B_2 that infect points in B_2^c in fact originate from points in $B_2 - B_1$, and we have control over the positions $U_1^1, \ldots U_{n_1}^1, U_1^2, \ldots, U_{n_2}^2$ of these points (the upper index indicating the infection type that infect these points). Then it suffices to couple the infection evolution of Z and Z' after time T ("final evolution"), and to construct an infection evolution of Z' such that at time T all points U_j^i are i-infected and no point of B_2^c is infected at all ("initial evolution"). In [DHB] this initial infection evolution of Z' was constructed by letting both infection types grow by sequences of very small infection outbursts that let both types grow close to some lines connecting the initial configuration with the corresponding points U_j^i (making use of (1.3)). Without condition (1.3) this approach seems to be hopeless, as it is extremely difficult to 1-infect a point U_j^1 but none of the points U_k^2, when there are a lot of points U_k^2 very close to U_j^1 and we are only allowed to use infection outbursts whose size is bounded away from 0. In the proof of the corresponding result in the discrete model in [DH2] there is a similar problem, which is solved by focusing on a special point $U_{j_0}^1$ and constructing an initial evolution infecting the point $U_{j_0}^1$ with type 1, but all points U_k^2 with type 2 before time T (assuming that $\beta_1 \geq \beta_2$). In order to be able to make use of the coupled infection evolution after time T, the point $U_{j_0}^1$ has to be chosen carefully, and there has to be a way of making sure that there are no infection outbursts from points we do not have any control of. The main reason that these unwanted outbursts can be prevented is that in the discrete model infections always originate from the boundary of the region that is already infected, which of course is not the case in the continuum model. Our strategy to prove Theorem 1 will be to adapt the trick from the lattice model, to enable us to construct an initial evolution. Due to the difficulties described above this will still be nontrivial. Additionally we need a new technique to prevent unwanted outbursts in a given region. One way to control outbursts in a special region is to make this region completely immune to infection. Thus one of our key tools will be a slightly different version of the CR-model that has immune regions and (for technical reasons) also allows delayed infections from the initial configuration, which will be called CR-model with generalized initial configuration.

A generalized initial configuration of an n-type CR-model will be of the form $\Gamma = (\Gamma_0, \Gamma_1, \ldots, \Gamma_n)$, where for $i \neq 0$ $\Gamma_i = (\Gamma_{i,t})_{t \geq 0}$ such that $\Gamma_{i,t} \neq \emptyset$ only for a finite number of values of t, and such that Γ_0 and $\Gamma_{i,t}$ $(1 \leq i \leq n, t \geq 0)$ are disjoint bounded Borel subsets of $\mathbb{R}^d$. Γ_0 can be thought of a set that never will be infected (immune region), and $\Gamma_{i,t}$ is immune up to time t, and then gets i-infected (delayed initial configuration). We will use the shorthand notation $\Gamma_\cup := \Gamma_0 \cup \bigcup_{i,t} \Gamma_{i,t}$. Additionally we are given growth parameters $\beta = (\beta_1, \ldots, \beta_n)$ and a radius distribution ρ as before. The corresponding stochastic process will again be denoted by $Z = (Z_t^i)_{i \in \{1,\ldots,n\}, t \geq 0}$, where Z_t^i denotes the i-infected subset of $\mathbb{R}^d$ at time $t \geq 0$. The dynamics of outbursts is a generalization of that for the CR-model with standard initial configurations. It is defined by the following properties:

- Z is a Markov process (which is time-homogeneous iff there is no time delay in the initial configuration, i.e. iff $\Gamma_{i,t} = \emptyset$ for $t > 0$).
- In $\Gamma_\cup$ we have a deterministic infection evolution as described by the initial configuration Γ.
- Regions of $\Gamma_\cup^c$ can only be infected by ball shaped infection outbursts. If the center of such a ball is i-infected, then the outburst i-infects all points within the ball that are not yet infected and not in $\Gamma_\cup$.
- At a given time t an outbursts of type i occurs at rate $\beta_i \lambda^d(Z_t^i)$. The center of the corresponding ball is chosen uniformly in Z_t^i and the radius is chosen w.r.t. ρ.

The above properties uniquely define the distribution of Z. We note that the CR-model with standard initial configuration is a special case of the CR-model with generalized initial configuration $\Gamma = (\Gamma_0, \ldots, \Gamma_n)$, corresponding to the case $\Gamma_0 = \emptyset$ and $\Gamma_{i,t} = \emptyset$ for $t > 0$. In Section 2 we will construct the CR-model with generalized initial configuration from a Poisson point process. What we actually need in our proof of Theorem 1 is a shape theorem for CR-models with generalized initial configurations, see Section 2.4.

This article is organized as follows: In Section 2 we clarify some notation, describe two ways to construct a CR-model from a Poisson point process, and state some important properties needed later. The proof of the corresponding lemmas and theorems is relegated to Section 3. In Section 4 we prove Theorem 1. The proof of the corresponding lemmas is relegated to Setion 5.

2 Setting

2.1 The underlying space and point process

First we will introduce some notation concerning subsets of $\mathbb{R}^d$ and point processes. Let $d(x,y) := |x-y|$ be the Euclidean distance between two points $x, y \in \mathbb{R}^d$. We will use $d(.,.)$ in the usual way also for the distance of a point and a set or the distance of two sets. For a subset $A \subset \mathbb{R}^d$ and a given $r > 0$ let

$$A_{+r} = \{x \in \mathbb{R}^d : d(x,A) \leq r\}$$

the r-enlargement of A. A_{+r} always is a closed set. For $x \in \mathbb{R}^d$, $r > 0$ let $B(x,r) := \{x\}_{+r}$ denote the closed ball with center x and radius r. On $\mathbb{R}^d$ we consider the Borel-σ-algebra denoted by $\mathcal{B}^d$, and the Lebesgue measure on $(\mathbb{R}^d, \mathcal{B}^d)$ denoted by λ^d. Let $(\mathbb{R}_+, \mathcal{B}_+, \lambda_+)$ be the restriction of $(\mathbb{R}, \mathcal{B}, \lambda)$ to $\mathbb{R}_+ = [0, \infty)$. The support of a set $A \in \mathcal{B}^d$ will be denoted by

$$\operatorname{supp}(A) := \{x \in \mathbb{R}^d : \lambda(A \cap B(x,r)) > 0 \text{ for all } r > 0\}.$$

If $A \subset \operatorname{supp}(A)$ then $\operatorname{supp}(A) = \overline{A}$, where $\overline{A}$ is the closure of A.

Considering point processes we will be mainly concerned with points $p = (x, s, r, w) \in \mathbb{R}^d \times \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+ =: \mathbb{R}^d_\times$. We will refer to the components of such a point p as position x, time s, radius r and strength w. For $A \subset \mathbb{R}^d$ and time interval $I \subset \mathbb{R}_+$ by abuse of notation we will sometimes consider A and $A \times I$ as subsets of $\mathbb{R}^d_\times$. We denote the Borel-σ-algebra on $\mathbb{R}^d_\times$ by $\mathcal{B}^d_\times$. A point configuration X is defined to be a subset of $\mathbb{R}^d_\times$ that is locally finite, i.e. $X \cap B(0,r) \times [0,T] \times \mathbb{R}_+ \times [0, \beta]$ is finite for all $r \geq 0$, $T \geq 0$, and $\beta \geq 0$. Let $\mathcal{X}$ denote the set of all point configurations. For $E \in \mathcal{B}^d_\times$ and $X \in \mathcal{X}$ let $N_E(X)$ be the number of points of X in E, and $\mathcal{F}_E$ the σ-algebra on $\mathcal{X}$ generated by all counting variables $N_{E'}$, where $E' \in \mathcal{B}^d_\times$ such that $E' \subset E$. $\mathcal{F} := \mathcal{F}_{\mathbb{R}^d_\times}$ will be the σ-algebra usually associated to $\mathcal{X}$. Sets $M \in \mathcal{F}_E$ will be called events depending only on (the information in) E.

For a given radius distribution ρ, which is a probability measure on $(\mathbb{R}_+, \mathcal{B}_+)$, we consider the Poisson point process with intensity measure $\lambda^d_\times := \lambda^d \times \lambda_+ \times \rho \times \lambda_+$ and denote its distribution by $\mathbb{P}$. $\mathbb{P}$ is the unique probability measure on $(\mathcal{X}, \mathcal{F})$ such that for bounded $E \in \mathcal{B}^d_\times$ the counting variable N_E is a Poisson random variable with parameter $\lambda^d_\times(E)$, and for disjoint sets $E_1, E_2, \ldots \in \mathcal{B}^d_\times$ the σ-algebras $\mathcal{F}_{E_1}, \mathcal{F}_{E_2}, \ldots$ are independent.

2.2 Constructing the model from a Poisson point process

Many proofs rely on the comparison of different continuum Richardson models (CR-models), so we have to construct certain couplings between CR-models. All of these couplings rely on the construction of a CR-model from a Poisson point process. In fact, we will give two methods to do this for the more general case of an n-type CR-model in d dimensions with growth rates $(\beta_1, \ldots, \beta_n)$

($\beta_i \geq 0$), radius distribution ρ satisfying (1.1), and generalized initial configuration $\Gamma = (\Gamma_0, \Gamma_1, \ldots, \Gamma_n)$, where for $i \geq 1$ $\Gamma_i = (\Gamma_{i,t})_{t \geq 0}$ such that $\Gamma_{i,t} \neq \emptyset$ only for a finite number of values of t, and such that Γ_0 and $\Gamma_{i,t}$ ($1 \leq i \leq n, t \geq 0$) are disjoint bounded Borel subsets of $\mathbb{R}^d$. The CR-model will be constructed from a Poisson point process $\mathbb{P}$ with intensity measure $\lambda^d_\times := \lambda^d \times \lambda_+ \times \rho \times \lambda_+$ on $(\mathbb{R}^d_\times, \mathcal{B}^d_\times)$. The CR-model will be denoted by $Z = (Z_t)_{t \geq 0}$, where $Z_t = (Z^1_t, \ldots, Z^n_t)$, and we think of Z^i_t as the subset of $\mathbb{R}^d$ that is infected with type i at time t. We will use the shorthand notation $Z^\cup_t := Z^1_t \cup \ldots \cup Z^n_t$. Let us first consider a point configuration $X \in \mathcal{X}$, and construct a deterministic infection evolution $Z = Z(X)$ for this X (but we should think of X as a typical configuration produced by $\mathbb{P}$).

Before we give a rigorous description of how to construct an infection evolution from a point configuration X in two different ways, we will describe these methods by more intuitive pictures. The first method, which we will call "scanning for points starting at time of infection", can be described as follows: No point of Γ_0 will ever get infected. Whenever $\Gamma_{i,t} \neq \emptyset$, this region is i-infected at time t and immune before. Whenever a new region A is i-infected at some time t, it gets a scanning device that starts scanning for points of X in $A \times [t, \infty) \times \mathbb{R}_+ \times [0, \beta_i]$ (so that a point of X with time $s \geq t$ will be scanned at time s). Whenever this device scans a point $p = (x, s, r, w)$, it produces an outburst that i-infects all points of $B(x, r)$ outside of $\Gamma_\cup$ that are not yet infected. This construction is changed only slightly in the second method, which we will call "scanning for points starting at time 0": Here the scanning device attached to a region A that gets i-infected at time t starts scanning for points of X in $A \times [0, \infty) \times \mathbb{R}_+ \times [0, \beta_i]$ (so that a point of X with time $s \geq 0$ will be scanned at time $t + s$). The rigorous description of these methods will be a bit technical because we have to consider the outbursts produced by the different scanning devices in their temporal order. Both methods are based on the following recursive construction:

At time $t = 0$ we define $Z^i_0 = \Gamma_{i,0}$. Now suppose that we have constructed an infection evolution $(Z_t)_{0 \leq t \leq t'}$ up to some $t' \geq 0$. Let $t^{in} := \min\{s > t' : \Gamma_{i,s} \neq \emptyset \text{ for some } i\}$ be the next time of an infection from the initial configuration, where we define $\min \emptyset = \infty$. By scanning the underlying point configuration X (as specified below) we may find a point $p = (x_p, s_p, r_p, w_p) \in X$, a corresponding type i_p and time $t_p \in (t', t^{in})$. If the scanning of X does not produce this information, then we say that $\Gamma_{i,t^{in}}$ is i-infected at time t^{in} and define $Z^i_t := Z^i_{t'} \cup \Gamma_{i,t'}$ for all $t \in (t', t^{in}]$ and $i \in \{1, \ldots, n\}$, which finishes the recursive step. If the scanning of X produces the above information, then we call $B_p := B(x_p, r_p)$ the (infection) outburst of type i_p produced by the point p at time t_p. The region $A_p := B_p - \Gamma_\cup - Z^\cup_{t'}$ is said to be i_p-infected at time t_p by the corresponding outburst. If $A_p \neq \emptyset$ the outburst is said to be effective. For all $t \in (t', t_p]$ we define $Z^i_t := Z^i_{t'}$, the only exception being that for $(t, i) = (t_p, i_p)$ we define $Z^{i_p}_{t_p} := Z^{i_p}_{t'} \cup A_p$, which finishes the recursive step.

Using a construction of this type we know that if we have constructed the infection evolution up to some time t', we can extend the construction to some interval $(t', t'']$ of positive length. We call the configuration X good, if the lengths of these intervals don't add up to something finite and if nothing goes wrong while scanning X, so that for a good configuration X indeed Z_t is well defined for all times $t \geq 0$. As in the above construction infected regions are a union of bounded sets, we know that Z_t^i is always bounded. By construction the sets Z_t^i $(1 \leq i \leq n)$ are disjoint for every t, and the sets Z_t^i are increasing in t for every i. A time t is called a time of growth if $Z_s^{\cup} \subsetneq Z_t^{\cup}$ for all $s < t$, i.e. if $t = t_p$ for some effective outburst p or $\Gamma_{i,t} \neq \emptyset$ for some i. Many proofs of assertions about an infection evolution are by induction on times of growth, i.e. induction on n, where $(t_n)_{n \geq 0}$ is the ordered sequence of all times of growth. Finally we call a sequence $(p_n)_{n \geq 1}$ of points $p_n \in X$ an i-infection path if every p_n produces an effective outburst that i-infects the position of p_{n+1}.

For "scanning X starting at time of infection" let t' be a given time such that $(Z_t)_{0 \leq t \leq t'}$ is a known infection evolution up to time t', and let t^{in} be the first time after t' that we have an infection by the initial configuration. Consider the set

$$\{(x, s, r, w) \in X : \exists i : x \in Z_{t'}^i, s \in (t', t^{in}], \text{ and } w \leq \beta_i\}, \tag{2.1}$$

and assume that the points of this subset of X are ordered w.r.t. their times s. If the set in (2.1) is empty, we say that the scanning doesn't produce a point. Otherwise, define $p = (x_p, s_p, r_p, w_p) \in X$ to be its minimum, i_p the unique type such that $x_p \in Z_{t'}^{i_p}$ and $t_p := s_p$. If the minimum is not unique or $t_p = t^{in}$ we say that the scanning goes wrong. (These cases could easily be taken care of, but that would make the construction more complicated.) For a good configuration X this construction is in accordance with the intuitive picture for this method described above. The important feature of this type of scanning is that for a good configuration X every point $p = (x, s, r, w) \in X$ such that x gets i-infected before time s and $w \leq \beta_i$ for some i produces an outburst, at time s, and other points don't produce outbursts.

For "scanning X starting at time 0" let again t' be a given time such that $(Z_t)_{0 \leq t \leq t'}$ is a known infection evolution up to time t', and let t^{in} be the first time after t' that we have an infection by the initial configuration. Let $\mathcal{A}(t')$ be the set of all triples (A, t, i) $(A \in \mathcal{B}^d$, $0 \leq t \leq t'$, $i \in \{1, \ldots, n\})$ such that (in the construction of the infection evolution up to time t') at some time t the set A was i-infected, either by an effective outburst or by the initial configuration. Consider the set

$$\bigcup_{(A,t,i) \in \mathcal{A}(t')} \{t + p : p = (x, s, r, w) \in X, x \in A, t + s \in (t', t^{in}], \text{ and } w \leq \beta_i\}, \tag{2.2}$$

where $t + (x, s, r, w) = (x, t + s, r, w)$, and assume that the points of this set are ordered w.r.t. to their corresponding times. If the set in (2.2) is empty, we say that the scanning doesn't produce a point. Otherwise, let $t+p$ be its minimum, where $p=(x_p,s_p,r_p,w_p)\in X$. There is a unique triple (A_p,i_p,t_p) such that $x_p\in A_p$, so this defines i_p and t_p. If the above minimum is not unique or $t_p=t^{in}$ we say that the scanning goes wrong. Again, for good X this construction is in accordance with the intuitive picture provided above. The important feature of this construction is that for a good configuration X a point $p=(x,s,r,w)\in X$ such that x gets i-infected at time t produces an outburst at time $t+s$ if $w\le\beta_i$ for some i, and doesn't produce an outburst otherwise.

The above constructions are deterministic, but choosing X according to the Poisson point process $\mathbb{P}$, both constructions give a random infection evolution defining the CR-model. As in [D], Prop. 2.1, for the usual one-type CR-model it can be shown that the configurations produced by $\mathbb{P}$ are good a.s. for both constructions, so that in both constructions Z_t^i a.s. is well defined for all $i\in\{1,\dots,n\}$ and $t\ge 0$ (this is a consequence of property (1.1) of the radius distribution ρ), and it is easy to check that in both constructions the process has the properties described in Section 1. The only instance where we will use "scanning from time 0" is in the proof of Theorem 4. In every other instance when a CR-model is related to a point process or a point configuration we will assume that the CR-model is constructed by "scanning from time of infection". Whenever we consider two or more initial configurations we write $Z_t^i(\Gamma)$ to indicate which initial configuration the process uses. In the special case of a standard initial configuration, the above constructions are slightly easier, see [D], [DHB] and [DH1].

2.3 Prescribing an infection evolution

For this subsection let Z denote a two-type CR-model with growth parameters $\beta_1,\beta_2>0$ and standard initial configuration $\Gamma=(\Gamma_1,\Gamma_2)$ with $\lambda^d(\Gamma_i)>0$. In the proof of Theorem 1 we will have to construct an infection evolution that has certain properties with positive probability. In order to have some control over possible radii produced by ρ, we choose a $d_0>0$ with the property

$$\rho([d_0,d_0+\epsilon])>0 \quad \text{for all } \epsilon>0. \tag{2.3}$$

For example, we can set $d_0:=\inf\{r>c:\rho([0,r])>\rho([0,c])\}$ for some small c with $\rho([0,c])<1$.

Lemma 1 (Constructing infection evolutions) *Let $d_0>0$ as above, $0<\delta<d_0/2$, $0\le T_1<T_2$, $B,C\subset\mathbb{R}^d$ bounded Borel sets and $c_0\in C$. Suppose that $C_{+d_0+2\delta}\subset B$ and C is of one of the following types:*

(a) $C=\{c_0,\dots,c_n\}$, where $(c_i)_i$ is a sequence of points such that for every $i\ge 1$ there is a $j<i$ with $d(c_j,c_i)\le d_0-2\delta$.

(b) C is a bounded, pathwise connected set.

For this choice of parameters there is an event M of point configurations such that M only depends on $B \times (T_1, T_2]$, $\mathbb{P}(M) > 0$, and for any CR-model Z we $\mathbb{P}$-a.s. have

$$\begin{aligned} &\{B(c_0, \delta) \subset Z^1_{T_1},\ Z^{\cup}_{T_1} \subset B,\ C_{+\delta} \cap Z^2_{T_1} = \emptyset\} \ \cap\ M \\ \subset \quad &\{C_{+d_0-2\delta} - Z^2_{T_1} \subset Z^1_{T_2} \subset C_{+d_0+2\delta} \cup Z^1_{T_1},\ Z^2_{T_2} = Z^2_{T_1}\}. \end{aligned} \tag{2.4}$$

Definition 1 (Growth along a set) *An event M with the properties given in Lemma 1 is said to describe the growth of type 1 along C (within B, starting at c_0) in the time interval $(T_1, T_2]$ using outbursts of size d_0 with precision δ.*

Definition 1 is of course motivated by property (2.4), which roughly says that whenever at time T_1 a neighborhood of c_0 is 1-infected and C is not 2-infected, the infection evolution caused by M ensures that in the time interval $(T_1, T_2]$ type 1 infects C_{+d_0} (as far as that is possible), whereas type 2 doesn't grow at all. We note that Lemma 1 and Definition 1 apply analogously to the growth of type 2 along a set; for this the roles of type 1 and 2 have to be interchanged. An important special case of Lemma 1, case (b), is that C is the trace of some continuous curve $\gamma : [0, 1] \to \mathbb{R}^d$ with $c_0 = \gamma(0)$.

2.4 Some properties of the model

For a given CR-model with generalized initial configuration and a Borel set B let τ_B denote the first time all points of B are infected, and ζ_B denote the time of the last effective outburst generated by a point with position in B. These random variables may take the value ∞, but we will show that this is not the case a.s.. Let us first consider a CR-model with standard initial configuration.

Lemma 2 *Let Z be a d-dimensional n-type CR-model with standard initial configuration $\Gamma = (\Gamma_1, \ldots, \Gamma_n)$ such that $\lambda^d(\Gamma_i) > 0$, growth rates $\beta_1, \ldots, \beta_n > 0$ and radius distribution ρ satisfying* (1.1)*. Let $B \subset \mathbb{R}^d$ be a bounded Borel set.*

- *(a) B is completely infected w.r.t. Z after a finite time, i.e. $\tau_B < \infty$ a.s..*
- *(b) There are only finitely many effective infection outbursts w.r.t. Z produced by points with position in B, i.e. $\zeta_B < \infty$ a.s..*

Very similar results have been obtained before. Note that in (b) we are not assuming the strong condition (1.2) on the radius distribution (compare to Lemma 4.5(a) in [DHB]). The proof of part (b) basically relies on the shape theorem stated below. For its proof we refer to [DHB].

Theorem 3 *Let Z be a d-dimensional one-type CR-model with standard initial configuration Γ with $\lambda^d(\Gamma) > 0$, growth rate $\beta > 0$ and radius distribution ρ satisfying* (1.2)*. There is a real $\mu > 0$ such that for all $0 < \epsilon < 1$ we almost surely have*

$$(1 - \epsilon) B(0, \beta\mu^{-1}) \subset Z_t/t \subset (1 + \epsilon) B(0, \beta\mu^{-1}) \quad \textit{for sufficiently large } t.$$

μ is independent of β and Γ.

We will need a slightly stronger version of Lemma 2 for a CR-model $\tilde{Z}$ with generalized initial configuration $\tilde{\Gamma}$, but we have to be careful as there may be regions that never will be infected since they are enclosed by a thick layer of the immune region $\tilde{\Gamma}_0$. Let $B_{\tilde{\Gamma}}$ be the smallest ball enclosing $\tilde{\Gamma}_{\cup}$ centered at the origin, and let $L := \{\exists t \geq 0 : \tilde{Z}_t \cap B_{\tilde{\Gamma}}^c \neq \emptyset\}$ be the event that the infection leaves $B_{\tilde{\Gamma}}$. We consider a point configuration X and the corresponding infection evolution $\tilde{Z}$ constructed from X by scanning from time of infection, and say that a point $p = (x, s, r, w) \in X$ has enough power to effectively infect points outside of $B_{\tilde{\Gamma}}$ if $B(x, r) \cap (\tilde{Z}_s^{\cup} \cup B_{\tilde{\Gamma}})^c \neq \emptyset$ (but x is not necessarily infected at time s, so p might not produce any outburst). Let $\tilde{\zeta}_B$ be the time of the last point of the underlying Poisson process with position in B that has enough power to effectively infect points outside of $B_{\tilde{\Gamma}}$.

Lemma 3 *Let $\tilde{Z}$ be a d-dimensional n-type CR-Model with generalized initial configuration $\tilde{\Gamma} = (\tilde{\Gamma}_0, \ldots, \tilde{\Gamma}_n)$, growth rates $\beta_1, \ldots, \beta_n > 0$ and radius distribution ρ. Let $B \subset \mathbb{R}^d$ be a bounded Borel set. On L we have a.s.:*

- *(a) $\tau_{B - B_{\tilde{\Gamma}}} < \infty$, i.e. $B - B_{\tilde{\Gamma}}$ is completely infected after a finite time.*
- *(b) $\tilde{\zeta}_B < \infty$, i.e. there are only finitely many points in B that have enough power to effectively infect points outside of $B_{\tilde{\Gamma}}$.*

We note that in (a) we considered $B - B_{\tilde{\Gamma}}$ rather than B so that we wouldn't have to worry about points enclosed by a thick layer of $\tilde{\Gamma}_0$. Again part (b) relies on a corresponding version of the shape theorem:

Theorem 4 *Let $\tilde{Z}$ be a d-dimensional one-type CR-model with generalized initial configuration $\tilde{\Gamma}$, growth rate $\beta > 0$ and radius distribution ρ satisfying (1.2). There is a real $\mu > 0$ such that for all $0 < \epsilon < 1$ on L we almost surely have*

$$(1 - \epsilon) B(0, \beta\mu^{-1}) \subset (\tilde{Z}_t \cup B_{\tilde{\Gamma}})/t \subset (1 + \epsilon) B(0, \beta\mu^{-1}) \quad \text{for sufficiently large } t.$$

μ is independent of β and $\tilde{\Gamma}$.

3 Proof of the results from the last section

For the following proofs we set $\beta_\wedge := \min_{1\leq i\leq n} \beta_i$ and $\beta_\vee := \max_{1\leq i\leq n} \beta_i$.

3.1 Constructing infection evolutions: Lemma 1

For the proof of Lemma 1, case (a), we simply describe a finite sequence of outbursts: Let $t_i := T_1 + \frac{i}{n+1}(T_2 - T_1)$ $(0 \leq i \leq n+1)$ and $B_i := B(c_i, \delta)$ $(0 \leq i \leq n)$. Let M be the set of all point configurations with exactly one point in each of $B_i \times (t_i, t_{i+1}] \times [d_0, d_0 + \delta) \times [0, \beta_1]$ for all $0 \leq i \leq n$, and no other points apart from these in $B \times (T_1, T_2] \times [0, \beta_\vee]$. It is easy to check that M has the desired properties.
We will prove Lemma 1, case (b), by reducing the situation described here to the situation of case (a). The compact set $C_{+d_0-2\delta}$ is covered by the union of all open balls with center in C and radius $d_0 - \delta$. We choose a finite subcover consisting of balls with centers $c_1, \ldots, c_n$. W.l.o.g. we may assume that $d(c_{i-1}, c_i) \leq d_0 - 2\delta$ for $1 \leq i \leq n$ (by joining the points c_i by continuous curves $\subset C$ and adding points of these curves that are sufficiently close to each other to this list of centers). Let M be chosen for $C' = \{c_0, \ldots, c_n\}$ according to part (a). By construction we have $\bigcup_i B(c_i, d_0 - \delta) \supset C_{+d_0-2\delta}$ and because of $c_i \in C$ we have $\bigcup_i B(c_i, d_0 + 2\delta) \subset C_{+d_0+2\delta}$ and $\bigcup_i B(c_i, \delta) \subset C_{+\delta}$. Thus the infection evolution given by M has the desired properties.

3.2 Standard initial configuration: Lemma 2

For the proof of part (a) we may assume that B is a ball with $B \supset \Gamma$ (by replacing B with a sufficiently big ball). Choose d_0 as in (2.3), and $\delta < d_0/2$. As in the proof of Lemma 1, case (b), we can choose $c_1, \ldots, c_n \in B$ such that $d(c_{i-1}, c_i) \leq d_0 - 2\delta$ for $1 \leq i \leq n$ and $\bigcup_i B(c_i, d_0 - \delta) \supset B$. We set $C_0 = \Gamma$ and $C_i = B(c_i, \delta)$, and for given $T > 0$ we set $t_i := \frac{i}{n+1}T$ $(0 \leq i \leq n+1)$. Let M_T be the set of all point configurations with at least one point in each of $C_i \times (t_i, t_{i+1}] \times [d_0, \infty) \times [0, \beta_\wedge]$ for all $0 \leq i \leq n$. On M_T B is completely infected at time T, and for $T \to \infty$ we have $\mathbb{P}(M_T) \to 1$. So by σ-continuity of $\mathbb{P}$, B is completely infected after a finite time a.s..

For the proof of part (b) we first show that there is a constant $v > 0$ s.t.

$$M_T := \{B(0, 2tv) \subset Z_t^\cup \text{ for all } t \geq T\} \uparrow \mathcal{X} \text{ a.s.} \quad \text{for } T \to \infty. \tag{3.1}$$

This basically is a consequence of a comparison argument and the shape theorem (Theorem 3): Let $f : [0, \infty) \to [0, 1)$ be a bijective increasing function such that $f(x) \leq x$, e.g. $f(x) = x/(1 + x)$. $\tilde{\rho} := \rho \circ f^{-1}$ has bounded support and thus satisfies (1.2). Let $\tilde{Z}$ be the one-type CR-model with initial configuration $\Gamma_\cup$, growth rate $\beta_\wedge > 0$ and radius distribution $\tilde{\rho}$. Theorem 3 applies to $\tilde{Z}$, which gives (3.1) for $\tilde{Z}$ instead of Z. Thus it suffices to construct a coupling of Z and $\tilde{Z}$ such that $\tilde{Z}_t \subset Z_t^\cup$ for all $t \geq 0$ a.s.. Let Z be constructed from X, and $\tilde{Z}$

from $\tilde{X} := \{(x, s, f(r), w) : (x, s, r, w) \in X\}$, scanning from time of infection, where X is chosen w.r.t. $\mathbb{P}$. Using this coupling the desired inclusion easily follows by induction on the times of growth of $\tilde{Z}$. This shows (3.1). If T is chosen so large that $B \subset B(0, Tv)$, on M_T every point $p = (x, s, r, w)$ producing an effective outburst in B after time T has the property $r \geq 2sv - Tv \geq sv$ and $w \leq \beta_\vee$, i.e. $N^{\text{eff}}_{B,T} \cdot 1_{M_T} \leq N_{E(B)}$, where $N^{\text{eff}}_{B,T}$ denotes the number of effective outbursts w.r.t. Z in B after time T and

$$E(B) := \{(x, s, r, w) \in \mathbb{R}^d_\times : x \in B, r \geq sv, w \leq \beta_\vee\}.$$

The $\mathbb{P}$-expectation of $N_{E(B)}$ is

$$\lambda^d_\times(E(B)) = \int_B d\lambda^d(x) \int d\rho(r) \int_0^{\beta_\vee} d\lambda^1(w) \int_0^{r/v} d\lambda^1(s) = \tfrac{\beta_\vee \lambda^d(B)}{v} \int \rho(dr) r,$$

where the last term is finite by condition (1.1) on ρ. So we have

$$\mathbb{P}(\{N^{\text{eff}}_{B,0} = \infty\} \cap M_T) = \mathbb{P}(\{N^{\text{eff}}_{B,T} = \infty\} \cap M_T) \leq \mathbb{P}(N_{E(B)} = \infty) = 0,$$

i.e. on M_T the number of effective outbursts in B is finite a.s., and the result follows from (3.1).

3.3 Generalized initial configuration: Lemma 3

For the proof of (a) we may assume that B is some big ball centered at 0 minus the set $B_{\tilde{\Gamma}}$. Choose d_0 as in (2.3). For rational parameters $c_0 \in B$, $t_0 > 0$ and $\delta < d_0/2$ let $L^{t_0}_{c_0,\delta}$ be the event that $Z^{\cup}_{t_0} \cap B$ contains $B(c_0, \delta)$. As $\bigcup_{t_0,c_0,\delta} L^{t_0}_{c_0,\delta} = L$ it suffices to show that $\tau_B < \infty$ a.s. on $L^{t_0}_{c_0,\delta}$ for arbitrary t_0, c_0, δ as above. This follows using the same construction as in the proof of Lemma 2, part (a), using $C_0 := B(c_0, \delta)$.

The proof of part (b) is almost exactly the same as that of part (b) of Lemma 2. Here we let $N^{\text{peff}}_{B,T}$ be the number points in B that have enough power to effectively infect points outside of $B_{\tilde{\Gamma}}$ after time T w.r.t the considered CR-model $\tilde{Z}$. We can use a comparison argument and Theorem 4 to show that

$$M_T := \{B(0, 2tv) \subset \tilde{Z}_t \cup B_{\tilde{\Gamma}} \text{ for all } t \geq T\} \uparrow L \text{ a.s.} \quad \text{for } T \to \infty$$

for some $v > 0$, and $N^{\text{peff}}_{B,T} \cdot 1_{M_T} \leq N_{E(B)}$ gives $\mathbb{P}(\{N^{\text{peff}}_{B,0} = \infty\} \cap M_T) = 0$.

3.4 Generalized Shape Theorem: Theorem 4

The basic idea of the proof is to compare the CR-model $\tilde{Z}$ with generalized initial configuration $\tilde{\Gamma}$ to the CR-model Z with standard initial configuration $\Gamma := B_{\tilde{\Gamma}}$, and then to use the shape theorem for Z. In order to be able to compare these models, we will use the following coupling: Let $\tilde{Z}$ and Z scan the same underlying Poisson point process "starting at time 0", see Section 2. For given rational times $0 \le T_1 \le T_2$ we define

$$M_{T_1,T_2} := \{\zeta_\Gamma \le T_1, Z_{T_1} \subset \tilde{Z}_{T_2} \cup \Gamma\} \cap L,$$

where ζ_Γ is the time of the last effective outburst in Γ w.r.t. Z. By Lemmas 2 and 3 we have $\zeta_\Gamma < \infty$ and $\tilde{\tau}_{B-\Gamma} < \infty$ a.s. on L for any bounded Borel set B and Z_{T_1} is bounded a.s. for any T_1. Thus we see that $\bigcup_{T_1,T_2} M_{T_1,T_2} = L$, so it suffices to show the generalized shape theorem on M_{T_1,T_2}. By the shape theorem for Z we know that there is a $\mu > 0$ independent of $\Gamma, \tilde{\Gamma}, \beta$ such that

$$(1-\epsilon/2)B(0,\beta\mu^{-1}) \subset Z_t/t \subset (1+\epsilon/2)B(0,\beta\mu^{-1}) \quad \text{for sufficiently large } t.$$

In order to see that this implies

$$(1-\epsilon)B(0,\beta\mu^{-1}) \subset (\tilde{Z}_t \cup \Gamma)/t \subset (1+\epsilon)B(0,\beta\mu^{-1}) \quad \text{for sufficiently large } t$$

on M_{T_1,T_2}, we use that $\frac{t-T_2}{t} \ge \frac{1-\epsilon}{1-\epsilon/2}$ for sufficiently large t and

$$\tilde{Z}_t \cup \Gamma \subset Z_t \subset \tilde{Z}_{t+T_2} \cup \Gamma \quad \text{for all } t \ge 0 \quad \text{on } M_{T_1,T_2}. \tag{3.2}$$

For the proof of (3.2) we fix a point configuration X, and we note that the chosen coupling of $\tilde{Z}$ and Z implies that every $p = (x,s,r,w) \in X$ with $x \notin \Gamma$ and $w \le \beta$ produces an outburst in both models, and in both models the time between the infection of x and the time p produces the outburst is s.

For the inclusion $\tilde{Z}_t \cup \Gamma \subset Z_t$ we observe that $Z_t \supset \Gamma$ is trivial and $\tilde{Z}_t \subset Z_t$ can be shown by induction on the growth times of $\tilde{Z}$: For $t = 0$ the assertion is trivial, and if $t > 0$ is a growth time, then either this growth is from the initial configuration $\tilde{\Gamma}$ of $\tilde{Z}$ (and in this case $\tilde{Z}_t \subset Z_t$ is an immediate consequence of the induction hypothesis) or there is a Poisson point $p = (x,s,r,w)$ producing an outburst at time t w.r.t. $\tilde{Z}$, i.e. x was infected in $\tilde{Z}$ at some time $t' = t - s < t$. By induction hypothesis $\tilde{Z}_{t'} \subset Z_{t'}$, so w.r.t. Z x is already infected at time t', i.e. p produces an outburst w.r.t. Z no later than $t' + s = t$.

For $Z_t \subset \tilde{Z}_{t+T_2} \cup \Gamma$ we argue with induction on the growth times of Z: For $0 \le t \le T_1$ the assertion is trivial since $Z_{T_1} \subset \tilde{Z}_{T_2} \cup \Gamma$ on M_{T_1,T_2}. If $t > T_1$ is a time of growth w.r.t. Z, there is a point $p = (x,s,r,w)$ producing an effective outburst in Z at time t. x must have been infected w.r.t. Z at time $t' = t - s < t$. So $x \in Z_{t'} \subset \tilde{Z}_{t'+T_2} \cup \Gamma$, where we have used the induction hypothesis. On M_{T_1,T_2} we have $t > T_1 \ge \zeta_\Gamma$ and thus $x \notin \Gamma$, so the above implies $x \in \tilde{Z}_{t'+T_2}$. Thus the outburst caused by p w.r.t. $\tilde{Z}$ occurs no later than $(t' + T_2) + s = t + T_2$. Therefore, the outburst doesn't destroy the inclusion.

4 Proof of the main result: Theorem 1

4.1 Outline of the strategy

We denote the two given CR-models by $Z(\Gamma)$ and $Z(\Gamma')$. Assuming

$$\mathbb{P}(G(\Gamma)) > 0, \qquad \text{and} \qquad \mathbb{P}(L_i(\Gamma')) > 0 \;\text{ for } i = 1, 2, \tag{4.1}$$

it suffices to show $\mathbb{P}(G(\Gamma')) > 0$. (4.1) implies $\beta_1, \beta_2 > 0$ and w.l.o.g. we may assume $\beta_1 \geq \beta_2$. Our basic strategy to show $\mathbb{P}(G(\Gamma')) > 0$ in the next subsections will be to decompose the evolution of $Z(\Gamma)$ into an initial part and a final part separated by a space-dependent time horizon. This will be chosen such that with positive probability the initial evolution will infect certain fundamental regions before the time horizon, and in the final evolution outbursts in these fundamental regions will yield paths to infinity for both types after the time horizon. After that we will describe an initial evolution of $Z(\Gamma')$ infecting the fundamental regions before the time horizon and having positive probability. Coupling the final evolution of $Z(\Gamma')$ to the final evolution of $Z(\Gamma)$ it can be seen that we also have coexistence for the initial condition Γ', i.e. $\mathbb{P}(\Gamma') > 0$.

4.2 Escaping the initial configuration in $Z(\Gamma')$

By (4.1) the initial configuration Γ' allows each type to escape some ball with positive probability. However, what we need is that Γ' also allows both types to escape some large ball at the same time with positive probability. This will be shown using the concept of a minimal enclosing ball (MEB). A MEB of a set $K \subset \mathbb{R}^d$ is defined to be a closed ball containing K with minimal radius, and it is easy to see that every bounded set has a unique MEB. We are mainly interested in the MEB of $\text{supp}(\Gamma'_1 \cup \Gamma'_2)$. We note that w.l.o.g. we may assume $\Gamma'_i \subset \text{supp}(\Gamma'_i)$ since a.s. there never will be an outburst originating from $\Gamma'_i - \text{supp}(\Gamma'_i)$. So we have $\text{supp}(\Gamma'_i) = \overline{\Gamma'_i}$. In order to have some control over possible radii produced by ρ we choose $d_0 > 0$ as in (2.3). If the radius r of the MEB of $\text{supp}(\Gamma'_1 \cup \Gamma'_2) = \overline{\Gamma'_1} \cup \overline{\Gamma'_2}$ has the property $\rho([r, \infty)) > 0$, we can choose d_0 such that $d_0 \geq r$.

We also need some notation concerning line segments: For given points $x_1 \neq x_2$ let $l_{x_1 x_2}$ denote the straight line passing through these points. We identify $l_{x_1 x_2}$ with $\mathbb{R}$ such that $x_1 < x_2$ so that we can use the induced order on $l_{x_1 x_2}$ along with the corresponding interval notation for line segments.

Definition 2 *We say that an infection configuration (A_1, A_2) has escape corridors specified by $x_1, x_2 \in \mathbb{R}^d$, $r_0, \delta_0 > 0$ whenever $d(x_1, x_2) > d_0 + 4\delta_0$, $B(x_i, \delta_0) \subset A_i \subset B(0, r_0)$, $(-\infty, x_1]_{+\delta_0} \cap A_2 = \emptyset$ and $[x_2, \infty)_{+\delta_0} \cap A_1 = \emptyset$.*

Lemma 4 *We can choose a time $T'_0 \geq 0$ and parameters $x_1, x_2 \in \mathbb{R}^d$, $\delta_0 < d_0/8$ and $r_0 > 2d_0$ such that $\mathbb{P}(M'_0) > 0$, where M'_0 is the event that $Z_{T'_0}(\Gamma')$ has escape corridors specified by x_1, x_2, r_0, δ_0.*

We note that on M_0' we can use Lemma 1 to let the two species grow along $(-\infty, x_1]$ and $[x_2, \infty)$ within $B(0, r_0)$ after time T_0' with precision δ_0.

4.3 Time horizon and fundamental regions for $Z(\Gamma)$

In the following we will define subsets of $M_0 := G(\Gamma)$ by specifying certain parameters of the infection evolution of $Z(\Gamma)$. After that we will have to show that the parameters can be chosen in a way such that these subsets of M_0 have positive probability. In Step 1 we use the parameters r_0, T_0' chosen in Lemma 4.

Step 1: Let $r_2 > r_1 > r_0$ and $B_i := B(0, r_i)$. Let M_1 be the set of configurations of M_0 such that

- at time T_0' the infected region is contained in B_1,
- no outburst with position in B_1 infects anything outside of B_2.

Step 2: Let $d_1 > 0$. Let M_2 be the set of configurations of M_1 such that $2d_1$ is a lower bound on the distances of the positions of effective outbursts in B_2 to

- the lines passing through the position of another effective outburst in B_2 and the origin,
- the boundaries of infection balls of outbursts that infect regions in B_2,
- the boundaries of the balls B_1 and B_2.

Step 3: Let $0 < d_2 < d_1$, $D^1, D^2_1, \ldots, D^2_k$ balls of radius d_2 ("fundamental balls") and $T(D^1), T(D^2_1), \ldots, T(D^2_k) > 0$ times associated to the balls. Let M_3 be the set of configurations of M_2 such that

- D^1 contains the starting point of an infinite type-1 infection path that never returns to B_2,
- the effective outbursts of type 2 within $B_2 - B_1$ are located in the balls $D^2_1, \ldots, D^2_k$ ($k \geq 1$, one ball for each outburst),
- and at the time $T(.)$ corresponding to a fundamental ball the position of the corresponding outburst is infected, but the outburst has not yet occurred.

Lemma 5 *The above parameters r_1, r_2, d_1, d_2, $D^1, \ldots, D^2_k$, $T(D^1), \ldots, T(D^2_k)$ can be chosen so that we have $\mathbb{P}(M_i) > 0$ for $i = 1, 2, 3$, and furthermore*

$$r_1 > r_0 + 4d_0, \quad r_2 > r_1 + 4d_0, \quad d_1 < \tfrac{d_0}{4}, \quad d_2 < \min\{\tfrac{d_1}{10}, \tfrac{d_1^2}{8d_0}\}. \tag{4.2}$$

We now choose the parameters according to the above lemma, and consider them to be fixed for the rest of the proof. $M_{\text{in,fi}} := M_3$ is the desired specification of the (initial and final) infection evolution of $Z(\Gamma)$. We have $\mathbb{P}(M_{\text{in,fi}}) > 0$ by Lemma 5, and on $M_{\text{in,fi}}$ the following properties hold:

- All fundamental balls are contained in $B_2 - B_1$ and keep a distance of at least $2d_1 - d_2$ to the boundaries of this set, all type-2 fundamental balls keep a distance of at least $2d_1 - d_2$ to the line through the center of D^1 and the origin, and the distance between two fundamental balls is at least $2d_1 - 2d_2$. The times corresponding to the fundamental balls are $> T_0'$.
- Each fundamental ball is infected as a whole (so the type-1 fundamental ball is completely 1-infected and the type-2 fundamental balls are completely 2-infected), and each fundamental ball contains the position of exactly one effective outburst. A fundamental ball is infected at the associated time, but the effective outburst in this ball occurs later.
- Every 2-infection of a region in B_2^c originating in B_2 in fact originates in one of the type-2 fundamental balls, and the type-1 fundamental ball is the starting point of an infection path to infinity for type 1 such that this path never returns to B_2.

We choose the time horizon on the fundamental balls to be the times associated with these balls, and the time horizon on B_2^c to be $T(B_2^c) := 0$. We will define the time horizon on the rest of the space later.

4.4 Initial evolution of $Z(\Gamma')$

In this section we only consider the model with initial configuration Γ', so here we set $Z := Z(\Gamma')$. We would like to construct an initial evolution starting from Γ' that infects all fundamental balls before the time horizon. More precisely we define

$$T' := \min\{T(D^1), T(D_1^2), \ldots, T(D_k^2)\} > T_0'$$

and M'_{in} to be the set of all configurations such that for Z at time T' ball D^1 is 1-infected, all balls D_i^2 $(1 \leq i \leq k)$ are 2-infected, and no point of B_2^c is infected, i.e.

$$M'_{\text{in}} := \{D^1 \subset Z^1_{T'}, \forall i : D_i^2 \subset Z^2_{T'}, Z_{T'} \subset B_2\}.$$

By definition M'_{in} only depends on the point configuration in

$$\Delta_{\text{in}} := B_2 \times [0, T'].$$

In order to show $\mathbb{P}(M'_{\text{in}}) > 0$ we give a step by step description of a suitable infection evolution having the desired properties. The main reason why this explicit construction is possible is that we only have to infect one small ball with type 1, but are allowed to infect everything else in B_2 with type 2. For the construction we choose fixed times $T_0' < T_1' < \ldots < T_4' := T'$, and we define a sequence of events M_i' describing a desirable infection state at time T_i'.

We will use interval notation for line segments of l_{0a}, where a denotes the center of D^1. Furthermore let c_1 denote the point on $[0, \infty) \subset l_{0a}$ with $d(c_1, 0) = \frac{r_0 + r_1}{2}$, let $c_2 = -c_1$ be its antipodal point, let $C := B_2 - B_1 - (\partial B_2)_{+d_0+d_2} -$



Figure 1: Sketch of the sets relevant in the definition of the initial evolution


T_1'
T_2'
T_3'
T_4'
■ = infected by species 1
▒ = infected by species 2
☐ = not infected
▓ = infected by species 1 or not at all
░ = infected by species 2 or not at all
▒ = infection state unspecified

Figure 2: Sketch of the infection states at time T_i' on the set M_i' for $1 \leq i \leq 4$.

$[0,\infty)_{+d_0+2d_2}$ and $C_+ := B_2 - B_1 - (\partial B_2)_{+d_1} - [0,\infty)_{+d_1}$, see Figure 1. The events M_i' (as shown in Figure 2) can now be defined by

$$
\begin{aligned}
&M_1' := \{B(c_1,d_2) \subset Z^1_{T_1'} \subset B_1 - [-r_1,c_2]_{+d_2}, B(c_2,d_2) \subset Z^2_{T_1'} \subset B_1 - [c_1,r_1]_{+d_2}\}\\
&M_2' := \{B(c_1,d_2) \subset Z^1_{T_1'} \subset B_1, C_{+d_0-d_2} - B_1 \subset Z^2_{T_2'} \subset B_2 - [c_1,r_2]_{+d_2}\}\\
&M_3' := \{D^1 \subset Z^1_{T_3'} \subset B_1 \cup [0,r_2]_{+d_1} \cap B_2, C_{+d_0-d_2} - B_1 \subset Z^2_{T_3'} \subset B_2\}\\
&M_4' := \{D^1 \subset Z^1_{T_4'} \subset B_2, C_+ \subset Z^2_{T_4'} \subset B_2\}.
\end{aligned}
$$

Lemma 6 *There are events* $M'_{(i-1)\to i}$ $(i = 1,2,3,4)$ *describing an infection evolution in* $(T'_{i-1}, T'_i]$ *such that* $\mathbb{P}(M'_{(i-1)\to i}) > 0$ *and* $M'_{i-1} \cap M'_{(i-1)\to i} \subset M'_i$.

Here M_0' is the event of positive probability chosen in Lemma 4. By the independence property of the Poisson point process we have $\mathbb{P}(M'_{i-1} \cap M'_{(i-1)\to i}) = \mathbb{P}(M'_{i-1})\mathbb{P}(M'_{(i-1)\to i})$, so Lemma 6 inductively gives $\mathbb{P}(M_i') > 0$ for $i = 0, \dots, 4$. As by construction all balls D_i^2 are contained in C_+, we have $M_4' \subset M'_{\text{in}}$, so the above implies that the local evolution in the Γ'-model has positive probability:

$$\mathbb{P}(M'_{\text{in}}) > 0 \tag{4.3}$$

4.5 Final evolution of $Z(\Gamma')$

Let us now consider the two-type CR-model $\tilde{Z} = Z(\tilde{\Gamma})$ with generalized initial configuration $\tilde{\Gamma} = (\tilde{\Gamma}_0, \tilde{\Gamma}_1, \tilde{\Gamma}_2)$, where

$$\tilde{\Gamma}_0 := D^0 := B_2 - (D^1 \cup D_1^2 \cup \dots D_k^2),$$

$\tilde{\Gamma}_{1,T(D^1)} := D^1$, $\tilde{\Gamma}_{2,T(D_1^2)} := D_1^2, \dots, \tilde{\Gamma}_{2,T(D_k^2)} := D_k^2$ and $\tilde{\Gamma}_{i,t} := \emptyset$ otherwise. We assume that $\tilde{Z}$ is constructed using the Poisson point process underlying $Z(\Gamma)$. On $M_{\text{in,fi}}$ the infection evolution of $\tilde{Z}$ in B_2^c is very similar to the one of $Z(\Gamma)$, differing only in that type 1 has been reduced considerably. However, we will show that in $\tilde{Z}$ type 1 is still strong enough to grow without bound. Let $M_{\text{fi}} := G(\tilde{\Gamma})$ denote the event that in $\tilde{Z}$ we have coexistence, i.e. both types grow without bounds.

Lemma 7 *We have* $M_{\textit{in,fi}} \subset M_{\textit{fi}}$ *and thus* $\mathbb{P}(M_{\textit{fi}}) > 0$.

We now would like to use M_{fi} for the final evolution of $Z(\Gamma')$, but unfortunately on $M'_{\text{in}} \cap M_{\text{fi}}$ we are not guaranteed to have coexistence for $Z(\Gamma')$. The problem is that in the construction of $\tilde{Z}$ from a Poisson point configuration all points in D^0 are ignored, so M_{fi} has no information on Poisson points in this region, whereas using M'_{in} for the initial evolution of $Z(\Gamma')$ it is very likely that D^0 will be at least partially infected at time T'. To guarantee coexistence in $Z(\Gamma')$, we thus would like to delete any point of the underlying Poisson process with position in D^0 that is strong enough to effectively infect points of B_2^c after time

T'. By Lemma 3 we know that $\tilde{\zeta}_{B_2} < \infty$ a.s. on M_{fi}, because $M_{\text{fi}} \subset L$. Thus we can choose a time $T'' > T'$ such that for

$$M'_{\text{fi}} := M_{\text{fi}} \cap \{\tilde{\zeta}_{B_2} < T''\} \quad \text{we still have} \quad \mathbb{P}(M'_{\text{fi}}) > 0. \tag{4.4}$$

M'_{fi} will be used for the final evolution in the Γ'-model, and setting $T(D^0) := T''$ we have defined a time horizon on all of $\mathbb{R}^d$. Since M'_{fi} is defined in terms of $\tilde{Z}$, it only depends on the part of the point configuration after the time horizon, i.e. on the point configuration in

$$\Delta_{\text{fi}} := \bigcup_{D \in \mathcal{D}} D \times [T(D), \infty), \quad \text{where} \quad \mathcal{D} = \{B_2^c, D^0, D^1, D_1^2, \ldots, D_k^2\}.$$

We now delete the remaining Poisson points that could interfere with the desirable final evolution. Let M'_{de} be the set of all point configurations without any point in

$$\Delta_{\text{de}} := \bigcup_{D \in \mathcal{D}} D \times [T', T(D)) \times \mathbb{R}_+ \times [0, \beta_1].$$

As $T(B_2^c) = 0$ we have $\lambda_\times^d(\Delta_{\text{de}}) < \infty$, and thus

$$\mathbb{P}(M'_{\text{de}}) > 0. \tag{4.5}$$

By construction the sets of Poisson configurations M'_{in}, M'_{de} and M'_{fi} depend on Δ_{in}, Δ_{de} and Δ_{fi} respectively, and these sets are disjoint. By the independence property of the Poisson point process we thus have

$$\mathbb{P}(M'_{\text{in,fi}}) = \mathbb{P}(M'_{\text{in}})\mathbb{P}(M'_{\text{de}})\mathbb{P}(M'_{\text{fi}}) > 0 \;\text{ for }\; M'_{\text{in,fi}} := M'_{\text{in}} \cap M'_{\text{de}} \cap M'_{\text{fi}},$$

where all probabilities are positive because of (4.3), (4.4) and (4.5). It is an immediate consequence of the construction that on $M'_{\text{in,fi}}$ the model $Z(\Gamma')$ has exactly the same infection evolution in B_2^c as the model $\tilde{Z}$, and we have coexistence for the model $\tilde{Z}$ on $M'_{\text{in,fi}}$ since $M'_{\text{in,fi}} \subset M'_{\text{fi}} \subset M_{\text{fi}}$. This implies that we also have coexistence for the model Z on $M'_{\text{in,fi}}$, i.e. $\mathbb{P}(G(\Gamma')) > 0$.

5 Proof of the Lemmas

5.1 Escape corridors: Lemma 4

Here we only consider the CR-model with initial configuration Γ', so let $Z := Z(\Gamma')$. Let $B = B(m,r)$ denote the MEB of $\overline{\Gamma'_1} \cup \overline{\Gamma'_2} = \mathrm{supp}(\Gamma'_1 \cup \Gamma'_2)$. We note that $r > 0$. We will make repeated use of the following property of MEBs:

$$\text{The MEB } B \text{ of a compact set } K \text{ is also the MEB of } \partial B \cap K. \tag{5.1}$$

We first show that we can choose an event M''_0 of positive probability such that on M''_0 at some time T the infection configuration Z_T contains distinct points $y_i \in \mathrm{supp}(Z^i_T)$ $(i = 1,2)$ such that using interval notation on $l_{y_1 y_2}$ we have

$$(-\infty, y_1 - d_0] \cap \overline{Z^2_T} = \emptyset \quad \text{and} \quad [y_2 + d_0, \infty) \cap \overline{Z^1_T} = \emptyset. \tag{5.2}$$

To show this we assume w.l.o.g. that $\partial B \cap \overline{\Gamma'_1} \neq \emptyset$ and $\partial B \cap \overline{\Gamma'_2} = \emptyset$. (By (5.1) one of these sets has to be nonempty. If it is the other way round we can argue similarly. If both are nonempty the assertion is trivial as we can simply choose $T = 0$ and $y_i \in \partial B \cap \overline{\Gamma'_i}$.) Thus by (5.1) B is the MEB of $\partial B \cap \overline{\Gamma'_1}$.

Case 1: $\rho([r,\infty)) > 0$. In this case we have $d_0 \geq r$. We choose $T = 0$ and $y_2 \in \overline{\Gamma'_2} - \{m\}$. As $B(m,r)$ is the MEB of $\partial B \cap \overline{\Gamma'_1}$, $B(y_2, r)$ doesn't enclose this set, so we can choose a $y_1 \in \partial B \cap \overline{\Gamma'_1}$ with $d(y_1, y_2) > r$. In order to see that y_1, y_2 have the above property, we only have to check that $[y_2 + d_0, \infty) \cap \overline{\Gamma'_1} \subset [y_2 + d_0, \infty) \cap B = \emptyset$, which follows from $\lambda^1(l_{y_1 y_2} \cap B - [y_1, y_2]) < 2r - r = r \leq d_0$.

Case 2: $\rho([r,\infty)) = 0$. Let M''_0 be the event that type 2 leaves B before the first type 1 outburst. To see that $\mathbb{P}(M''_0) > 0$ let $T'' > 0$, M''_1 the set of point configurations with no point in $\Gamma'_1 \times [0, T''] \times [0, \beta_1]$ and M''_2 the event that type 2 leaves B before time T''. We have $\mathbb{P}(M''_1) > 0$, and if T'' is chosen large enough we also have $\mathbb{P}(M''_2) > 0$. The event $M''_2|_{\Gamma_1'^c} := \{X \in \mathcal{X} : X|_{\Gamma_1'^c} \in M''_2\}$ is independent of M''_1 and we have $M''_2|_{\Gamma_1'^c} \supset M''_2 \supset M''_1 \cap M''_2|_{\Gamma_1'^c}$ as deleting outbursts of type 1 can only strengthen type 2. So

$$\begin{aligned} \mathbb{P}(M''_0) \geq \mathbb{P}(M''_1 \cap M''_2) \geq \mathbb{P}(M''_1 \cap M''_2|_{\Gamma_1'^c}) &= \mathbb{P}(M''_1)\mathbb{P}(M''_2|_{\Gamma_1'^c}) \\ &\geq \mathbb{P}(M''_1)\mathbb{P}(M''_2) > 0. \end{aligned}$$

On M''_0 at some time T an outburst $B' = B(x', r')$ of species 2 occurs that for the first time infects a region in B^c. As $r' < r$ a.s. and as B is the MEB of $\partial B \cap \overline{\Gamma'_1}$, B' can't enclose this set. So we can choose points $y_1 \in \partial B \cap \overline{\Gamma'_1} - B'$ and $y_2 \in \partial B \cap B'$. These points have the above property.

The infection configuration always is constant over a time interval of positive length, $\overline{Z^i_T}$ is always a compact set, the infected region is always bounded, and if the points y'_i are sufficiently close to y_i, these points still satisfy (5.2). This shows that on M_0'' there are rational parameters $T > 0$, $0 < \delta < d_0/8$, $y_1', y_2' \in \mathbb{R}^d$ and $r > 0$ such that (using interval notation on $l_{y_1'y_2'}$) we have

$$\begin{gathered}(-\infty, y_1' - d_0]_{+4\delta} \cap \overline{Z_T^2} = \emptyset,\ [y_2' + d_0, \infty)_{+4\delta} \cap \overline{Z_T^1} = \emptyset,\ d(y_1', y_2') > 4\delta, \\ Z_T^{\cup} \subset B(0, r) \ \text{ and } \ \lambda^d(B(y_i', \delta) \cap Z_T^i) > 0 \text{ for } i = 1, 2.\end{gathered} \tag{5.3}$$

Using the σ-additivity of $\mathbb{P}$ we thus can choose parameters as above such that the event M_{00}' that (5.3) holds for this particular choice of parameters has positive probability. Let M_{01}' be the set of all Poisson configurations with exactly one point in each of the sets

$$\big(B(y_i', \delta) \cap Z_T^i\big) \times (T, T+1] \times [d_0, d_0 + \delta) \times [0, \beta_2] \quad (i = 1, 2),$$

and no additional points in $B(0, r) \times (T, T+1] \times [0, \beta_1]$. The event $M_{01}' \in \mathcal{F}_{\mathbb{R}^d \times (T,T+1]}$ is independent of $M_{00}' \in \mathcal{F}_{\mathbb{R}^d \times [0,T]}$ and has positive probability. As a consequence $M_0' := M_{00}' \cap M_{01}'$ has positive probability. On M_0' we know that at time $T_0' := T + 1$ the infection configuration $Z_{T_0'}$ has escape corridors specified by the parameters $x_1 := y_1' - d_0 + 2\delta$ and $x_2 := y_2' + d_0 - 2\delta$, $r_0 := r + 2d_0$ and $\delta_0 := \delta$.

5.2 Specifying the evolution in the Γ-model: Lemma 5

Step 1: For any choice of $T > 0$ we have $M_0 \cap \{Z_{T_0'} \subset B(0, r_1)\} \cap \{\zeta_{B_1} < T\} \cap \{Z_T \subset B(0, r_2)\} \subset M_1$. The events $\{Z_{T_0'} \subset B(0, r_1)\}$, $\{\zeta_{B_1} < T\}$ and $\{Z_T \subset B(0, r_2)\}$ are increasing in r_1, T and r_2 respectively, and $\zeta_{B_1} < \infty$ a.s. by Lemma 2 and for fixed T Z_T is bounded a.s.. Thus we only have to choose first r_1, then T and then r_2 large enough in order to obtain $\mathbb{P}(M_1) > 0$.

Step 2: Observing that the event of all given distances being bounded from below by $2d_1$ is decreasing in d_1, we are left to show that the infimum of the distances is positive a.s.. Every single distance of those considered is positive a.s. by the properties of the Poisson point process, so it suffices to observe that the number of effective outbursts in B_2 is finite a.s. by Lemma 2 and the number of outbursts that infect a region in B_2 is finite a.s., as B_2 is completely infected after a finite time by Lemma 2. Therefore if we choose $d_1 > 0$ small enough, we have $\mathbb{P}(M_2) > 0$.

Step 3: We choose a small $d_2 > 0$ satisfying (4.2), and we fix a finite covering of B_2 consisting of balls of radius d_2. For any infection configuration it is possible to choose a finite number of balls with the properties described in Subsection 4.3, Step 3, and as in each ball the time span between the infection of the outburst position and the outburst at that position is positive a.s. we can choose a rational time serving as the time horizon on the ball. The number of ways to choose a finite set of fundamental balls from the finite covering and a rational time horizon is countable, so using the σ-additivity of $\mathbb{P}$ there is a specific set of fundamental balls and corresponding time horizons such that $\mathbb{P}(M_3) > 0$.

5.3 Local evolution in the Γ'-model: Lemma 6

$M'_{0\to 1}$: Infecting the balls $B(c_i, d_2)$. We will use the parameters chosen in Lemma 4 and $\delta := \delta_0$. Choosing two curves $C_i \subset B_{1/2} := B(0, r_0 + 2d_0)$ $(i = 1, 2)$ joining $x_i \in B_0$ and $c_i \in \partial B_{1/2}$ and choosing a time $T'_{1/2} \in (T'_0, T'_1)$, we can use Lemma 1 to let type 1 grow along C_1 in the time interval $(T'_0, T'_{1/2}]$ with precision δ and after that to let type 2 grow along C_2 in the time interval $(T'_{1/2}, T'_1]$ with precision δ. This infection evolution defines an event $M'_{0\to 1}$ with the desired properties provided that the curves C_i satisfy $d(C_1, Z^2_{T'_0}) > \delta$, $d(C_2, Z^1_{T'_0}) > \delta$, $d(C_1, C_2) = d(x_1, x_2) > d_0 + 4\delta$ and $d(C_1, B(c_2, d_2)) > d_0 + 2\delta$. For a possible construction of such curves see Figure 3. Here b_1 and b_2 are antipodal points on the surface of $B(0, r_0 + \frac{d_0}{2})$ such that $l_{b_1 b_2}$ is parallel to $l_{x_1 x_2}$. After passing b_i the curve C_i continues on the line in the hyperplane perpendicular to $l_{b_1 b_2}$ that hits $B_{1/2}$ in a point with minimal distance to c_i.



Figure 3: Construction of the curves C_1 and C_2

$M'_{1\to 2}$: Infecting most of $B_2 - B_1$ with type 2. We use Lemma 1 to let type 2 grow along the pathwise connected, bounded set $C' := C \cup [-r_1, c_2]$ in the time interval $(T'_1, T'_2]$ with precision $\delta := d_2/2$. This infection evolution defines an event $M'_{1\to 2}$ with the desired properties.

$M'_{2\to 3}$: Infecting D^1 with type 1. In this step we basically would like to let type 1 grow along $[c_1, r_2]$ with precision $\delta := d_2/2$ until it reaches D^1. In order to keep the 1-infections close to $[c_1, r_2]$, we have to ensure that for every outburst of type 1 all points of the corresponding ball that are too far away from $[c_1, r_2]$ already are 2-infected before the outburst occurs. In this case we say that species 2 provides sufficient containment for the outburst. In the following things get more complicated if D^1 is near ∂B_2, and thus outbursts infecting D^1 do not yet have sufficient containment. For a line segment l' of the line $l = l_{c_1 c_2}$ let $S_r(l')$ denote the cylindrical shell with axis l' and radius r, i.e. the set of all points x with $d(x, l) = r$ such that the projection of x onto l is on l'. In case of a single point x we write $S_r(x) := S_r(\{x\})$. In order to determine for which points $a' \in [c_1, a]$ an infection outburst at a' with precision δ has sufficient containment on M_2', we first note that

$$d(S_r(r_2 - s), \partial B_2) \geq s - r \quad \text{for all } 0 \leq r \leq s \leq r_2. \tag{5.4}$$

(This and other purely geometric relations will be proved at the end of this subsection.) (5.4) implies $S_{d_0+2d_2}([0, r_2 - 2d_0 - d_1]) \subset C \cup B_1$ and thus

$$S_{3d_2}([0, r_2 - 2d_0 - d_1]) \subset S_{d_0+2d_2}([0, r_2 - 2d_0 - d_1])_{+d_0-d_2} \subset C_{+d_0-d_2} \cup B_1 \tag{5.5}$$

On M_2' we know that $C_{+d_0-d_2} - B_1$ is 2-infected, so this shows that the above described infection outburst at a' has sufficient containment if $a' \in [c_1, a]$ is such that

$$a' \leq r_2 - 2d_0 - d_1 - (d_0 + 2\delta) = r_2 - (3d_0 + d_1 + d_2). \tag{5.6}$$

Thus, if $a' := a - (d_0 - 2\delta)$ satisfies (5.6) (i.e. if $a \leq r_2 - (2d_0 + d_1 + 2d_2)$) we can use Lemma 1 to let type 1 grow along the curve $[c_1, a']$ in the time interval $[T_2', T_3']$ with precision δ, and immediately get an event $M_{2\to3}'$ as desired. However, if $a > r_2 - (2d_0 + d_1 + 2d_2)$, we might need two more outbursts (of size d_0) to reach D^1 with containment not yet provided. Before we define a suitable infection evolution in this case, we will give an informal description, using Figure 4 which shows a halfplane starting from $l_{c_1 a}$. $a_0 < b_0 < a_1 < b_1 < a_2 < b_2 < a_3 = a$ are points chosen on $[c_1, a]$. The three points above $[c_1, a]$ indicate the points of intersection of the halfplane with the cylindrical shells $S_{d_1}(b_i)$. We would like to define $M_{2\to3}'$ to be the following infection evolution (where $T_2' = t_0' < t_1' < \ldots < t_5' = T_3'$):



Figure 4: Definition of the infection evolution of step $M_{2\to3}'$ in the difficult case. Some infection outbursts are indicated by segments of their boundaries, which are connected to their centers by dotted lines.

1. growth of type 1 along $[c_1, a_0]$ in $[t_0', t_1']$,
2. growth of type 2 along $S_{d_1}([b_0, b_1])$ in $[t_1', t_2']$,

3. growth of type 1 by a single outburst at a_1 in $[t'_2, t'_3]$,
4. growth of type 2 along $S_{d_1}([b_1, b_2])$ in $[t'_3, t'_4]$,
5. growth of type 1 by a single outburst at a_2 in $[t'_4, t'_5]$.

In every step the growth is meant to be within B_2 and with precision δ. (Note that in Steps 2 and 4 $S_{d_1}([b_i, b_{i+1}])$ consists of either one or two connected components, depending on whether $d > 2$ or $d = 2$. In the case of two connected components we let type 2 grow along each component.) The important part of the boundary of the infection outburst in every step is indicated in Figure 4. By Lemma 1 we have $\mathbb{P}(M'_{2\to 3}) > 0$, and Figure 4 shows that for $M'_2 \cap M'_{2\to 3} \subset M'_3$ we only have to ensure the following properties in the corresponding steps: In Step 1 we need sufficient containment, i.e. a_0 has to satisfy (5.6), and $B(a_1, d_2)$ has to be fully infected, i.e. for $i = 0$ we need

$$d(a_i, a_{i+1}) \leq d_0 - 2d_2. \tag{5.7}$$

For Step 2 we need

$$S_{d_1}(b_0)_{+d_2} \subset C_{+d_0-d_2}, \tag{5.8}$$

and in this step no part of $B(a_2, d_2)$ may be 2-infected, i.e. for $i = 1$ we need

$$d(S_{d_1}(b_i), a_{i+1}) \geq d_0 + 2d_2. \tag{5.9}$$

Also this step has to provide sufficient containment for the next, i.e. we need

$$B(a_i, d_0 + 2d_2) \subset [0, r_2]_{+d_1/2} \cup S_{d_1}([b_{i-1}, b_i])_{+(d_0-d_2)} \tag{5.10}$$

for $i = 1$. In Step 3 $B(a_2, d_2)$ has to be fully infected, i.e. we need (5.7) for $i = 1$. Steps 4 and 5 are completely analogous to Steps 2 and 3, so we just need (5.9), (5.10) and (5.7) for $i = 2$. Finally, to ensure that the growth is always within B_2 we need

$$S_{d_1}([b_0, b_2])_{+(d_0+d_2)} \cup B(a_2, d_0 + 2d_2) \subset B_2. \tag{5.11}$$

Defining $a_0 < a_1 < a_2 < a_3 := a$ and $b_i \leq a_{i+1}$ to be those points on $[c_1, a]$ satisfying

$$d(a_i, a_{i+1}) = d_0 - 2d_2 \quad \text{and} \quad d(S_{d_1}(b_i), a_{i+1}) = d_0 + 2d_2, \tag{5.12}$$

we can now check the above properties: (5.7) and (5.9) are trivially satisfied. It is easy to check that a_0 satisfies (5.6), using

$$a_3 = a \leq r_2 - 2d_1, \tag{5.13}$$

which is true by definition of a and d_1, and $8d_2 \leq d_1$. (5.8) follows from (5.5) provided $b_0 \leq r_2 - 2d_0 - d_1$, which follows from

$$a_i \leq b_i \leq a_i + d_1/2. \tag{5.14}$$



Figure 5: Illustrations for proofs of (5.4) and (5.14)

What remains to be checked are properties (5.4), (5.14), (5.10) and (5.11) . As can be seen from Figure 5, assertion (5.4) is equivalent to

$$\sqrt{(r_2 - s)^2 + r^2} + (s - r) \le r_2,$$

which holds for all $r \le s \le r_2$. Figure 5 also shows that (5.14) is equivalent to

$$0 \;\le\; d_0 - 2d_2 - \sqrt{(d_0 + 2d_2)^2 - d_1^2} \;\le\; d_1/2,$$

which can seen to be satisfied for $d_2 \le d_1^2/(8d_0)$ and $2d_1 \le d_0$. For (5.10) we observe that $[a_i - (b_i - a_i), a_i + (b_i - a_i)] \subset [b_{i-1}, b_i]$, which follows from (5.14), so by symmetry it suffices to check that



Figure 6: Illustration for proof of (5.10)

$$d(S_{d_1}(b_i + d_0 - d_2), a_i) \ge d(S_{d_1}(b_i), a_{i+1}) = d_0 + 2d_2,$$

see Figure 6. Here the equality follows from (5.12) and Figure 6 shows that the inequality follows from $a_i \le b_i$ and $b_i + d_0 - d_2 \ge a_i + d_0 - d_2 > a_{i+1}$. For (5.11) we first observe that $B(a_2, d_0 + 2d_2) \subset B_2$ follows from $r_2 - a_2 \ge 2d_1 + (d_0 - 2d_2) \ge d_0 + 2d_2$, where we have used (5.13). Furthermore we have

$$S_{d_1+d_2}(r_2 - d_0 - d_1 - 2d_2)_{+d_0+d_2} \subset B_2,$$

by (5.4), so it suffices to show that $b_2 \le r_2 - d_0 - d_1 - 2d_2$. This follows from (5.12), (5.13) and (5.14).

$M'_{3\to 4}$: Infecting the remaining part of $B_2 - B_1$ with type 2. Let $r'_2 := d(S_{d_1+d_2}(r_2-d_0-d_1-2d_2), 0)$, and $C' := B(0, r'_2) - B_1 - [0, r_2]_{+d_1+d_2}$. We define $M'_{3\to 4}$ by the growth of type 2 along C' in the time interval $[T'_3, T'_4]$ with precision $\delta := d_2/2$. By Lemma 1 we have $\mathbb{P}(M'_{3\to 4}) > 0$ and for $M'_3 \cap M'_{3\to 4} \subset M'_4$ it suffices to observe that on M'_3 no point of C'_{+d_2} is 1-infected, most of C' is 2-infected, by (5.4) we have $C'_{+d_0+d_2} \subset B_2$, and by choice of r'_2 we have $C'_{+d_0-d_2} \supset C_+$.

5.4 Restricting the strong type: Lemma 7

In this section we restrict our attention to the infection evolutions of $Z := Z(\Gamma)$ and $\tilde{Z}$ for a fixed point configuration $X \in M_{\mathrm{in,fi}}$. By definition of $M_{\mathrm{in,fi}}$, we know that Z^2 grows without bound and X contains an infinite type-1 infection path $(p_n)_{n\geq 1}$ w.r.t Z such that for $p_1 = (x_1, t_1, r_1, b_1)$ we have $x_1 \in D^1$ and $t_1 \geq T(D^1)$. It suffices to show that

- $\tilde{Z}^2_t \cap B_2^c \supset Z^2_t \cap B_2^c$ for all $t \geq 0$, and
- $(p_n)_n$ is an infinite type-1 infection path w.r.t $\tilde{Z}$.

To prove the first assertion we show that

$$\forall t \geq 0: \quad \tilde{Z}^{\cup}_t \subset Z^{\cup}_t, \tilde{Z}^1_t \subset Z^1_t, \text{ and } Z^2_t \cap B_2^c \subset \tilde{Z}^2_t \cap B_2^c$$

by induction on the times of growth of both models. At $t = 0$ the above is trivial, so suppose that $t > 0$ is a time of growth, and $t-$ is the previous time of growth. By induction hypothesis (IH) the assertion is true at time $t-$. As one of the models grows at time t, either $\tilde{\Gamma}_{i,t} \neq \emptyset$ for some i, or we have a point $p = (x, t, r, w) \in X$ producing an outburst in one of the models. In the first case the assertion is still true at time t as in the model Z every fundamental ball is infected at its time horizon. In the second case (IH) implies that p actually produces an outburst w.r.t. Z, so $x \in Z^{\cup}_{t-}$. If $x \in Z^1_{t-}$ the assertion is trivially true at time t. If $x \in Z^2_{t-}$ it suffices to check that

$$x \notin \tilde{Z}^1_{t-} \quad \text{ and } \quad (B(x,r) \cap B_2^c \neq \emptyset \;\Rightarrow\; x \in \tilde{Z}^2_{t-}),$$

because as a consequence the same infection outburst as in Z is produced in $\tilde{Z}$. The first assertion follows from $x \notin Z^1_{t-} = Z^1_t$ and (IH). For the second assertion assume that $B(x,r) \cap B_2^c \neq \emptyset$. If $x \in B_2$, then by construction x is in one of the type-2 fundamental balls and t is after the corresponding time horizon, which implies $x \in \tilde{Z}^2_{t-}$. If $x \notin B_2$, then $x \in Z^2_{t-} \cap B_2^c \subset \tilde{Z}^2_{t-} \cap B_2^c$. This concludes the induction proof.

To check that $(p_n)_n$ is an infinite type-1 infection path w.r.t $\tilde{Z}$, we suppose otherwise, i.e. there is a $k \geq 1$ such that $p_1, \ldots, p_k$ is a type-1 infection path w.r.t $\tilde{Z}$, but x_{k+1} is infected by an outburst $p = (x, t, r, w)$ different from p_k. As p_k still produces an outburst of infection 1 at time t_k, we have $t < t_k$. On the other hand we have $x \in \tilde{Z}^{\cup}_t \subset Z^{\cup}_t$, $\tilde{Z}^1_t \subset Z^1_t$ and the infection ball of p contains

x_{k+1}, so we know that x_{k+1} was infected w.r.t. Z at time t or before that. This is a contradiction as we know by construction that x_{k+1} was infected w.r.t. Z at time $t_k > t$.

Acknowledgments: We would like to thank H.-O. Georgii for bringing this topic to our attention and for helpful discussions and remarks.

References

[D] M. Deijfen, Asymptotic shape in a continuum growth model, Adv. Appl. Prob. 35 (2003), 303-318.

[DH1] M. Deijfen, O. Häggström, Coexistence in a two-type continuum growth model, Adv. Appl. Prob. 36 (2004), 973-980.

[DH2] M. Deijfen, O. Häggström, The initial configuration is irrelevant for the possibility of mutual unbounded growth in the two-type Richardson model, Comb. Prob. Computing 15 (2006), 345-353.

[DHB] M. Deijfen, O. Häggström, J. Bagley, A stochastic model for competing growth on $\mathbb{R}^d$, Markov Proc. Relat. Fields 10 (2004) 217-248.

[HP1] O. Häggström, R. Pemantle, First passage percolation and a model for competing spatial growth, J. Appl. Probab. 35 (1998), 683-692.

[HP2] O. Häggström, R. Pemantle, Absence of mutual unbounded growth for almost all parameter values in the two-type Richardson model, Stoch. Proc. Appl. 90 (2000), 207-222.

[R] D. Richardson, Random growth in a tessellation, Proc. Cambridge Phil. Soc. 74 (1973), 515-528.